

Annual Monitoring Report Summary
Downeast Lakes Land Trust,
Downeast Lakes Community Forest

2020

Prepared by Ernest Carle

Table of Contents

I. INTRODUCTION	3
II. ANNUAL MONITORING UPDATE	4
Timber Harvest	4
Other Forest Products.....	6
Unanticipated Removal or Loss	6
Regeneration.....	6
Focus Species Habitat Management Activities	7
DLLT Self-Designated Deer Management Areas.....	7
Grouse and Woodcock.....	8
Black Bear	10
Riparian Zone Management	10
Beaver.....	11
Brook Trout / Atlantic Salmon	11
Exotic and Invasive Plants.....	11
Harvest Impacts.....	14
Road Monitoring	16
Pesticides and Biological Control Agents	18
Social and Economic Monitoring	18
Common Loon Monitoring.....	18
Forest Inventory	20
Changes in Habitat Conditions	21
Deer Management Areas	22
American Marten.....	23
Grouse and Woodcock.....	23
Black-throated Blue Warbler/Mature Hardwood Forest.....	25
Hard Mast Management	25
Rare Species, Natural Communities, and other Special Habitats	26
Literature Cited.....	26

I. Introduction

Downeast Lakes Land Trust (DLLT) periodically monitors its forest to ensure that its management objectives for wildlife, recreation, timber production, and environmental protection are being met. Some items, such as the inventory of standing timber, are conducted periodically. Others, such as monitoring timber harvest operations, may be conducted on a weekly basis as operations are ongoing. This annual report summarizes the monitoring information for members of the land trust, members of the local community, and for others interested in the results of DLLT's management. For more information on DLLT's forest management, please contact the land trust or visit www.downeastlakes.org. The monitoring update includes annual summaries for the operating year. Periodic monitoring data that are updated every 5-10 years are included in Section III.

This report covers the Downeast Lakes Community Forest, which consists of the 27,080-acre Farm Cove Tract acquired by Downeast Lakes Land Trust in 2005 and expanded in 2008 by acquisition of the 6,628-acre Wabassus Lake Tract and in 2016 by the acquisition of the 21,853-acre West Grand Lake Tract. The management plan for all tracts was updated and adopted on March 2nd, 2015. No major changes in policy or management strategy have taken place.

II. Annual Monitoring Update

Timber Harvest

Timber Harvest Summary	2009-2015 (Average)		2016		2017		2018		2019		2020	
	Species	Prod.	Cords	Prod.	Cords	Prod.	Cords	Prod.	Cords	Prod.	Cords	Prod.
Hemlock	Stud	1473	Stud		Stud		Stud	52	Stud	1072	Stud	270
Hemlock	Pulp	1645	Pulp	222	Pulp	538	Pulp	1464	Pulp	2447	Pulp	1839
Hemlock	Logs	128	Logs	375	Logs	1003	Log	663	Log	685	Log	902
Hemlock	WTC	139	WTC	324	WTC	361	WTC					
Spruce	Logs	430	Logs	1	Logs							
Softwood ⁴	Pulp	562	Pulp	284	Pulp	103	Pulp	693	Pulp	3118	Pulp	1002
Spruce/Fir	Stud	233	Stud	1109	Stud	1970	Stud	2995	Stud	1928	Stud	1573
Red Pine	Land Run						Land Run	325				
White Pine	Logs	33	Logs	1273	Logs	243	Logs	366	Logs	356	Logs	1434
Hardwood	Pulp	1601	Pulp	2444	Pulp	2478	Pulp	4582	Pulp	2812	Pulp	2518
Hardwood	Logs	3						19		1		1
Hardwood	Veneer	1										
Hardwood	Firewood	14								43		33
Subtotal (without biomass)¹		6258		6032		6696		11157		12462		
MX Biomass ¹	Chips	1818	Chips	3136	Chips	4502	Chips	1311	Chips	230	Chips	0
Biomass ²	2013	1296										33
Biomass ³							2017	583				
Total w. biomass:		8267		9770		11781		12468		12692		9605

Monitoring Summary Report

¹ Biomass sales are typically incidental to planned harvest volumes and are composed of tops or limbs that are not considered within timber inventory. In 2009, 112 cords of biomass included in the figures above was harvested during maintenance of the Farm Cove Dam Road.

² Biomass in 2013 was also generated from a wildlife habitat project that had no commercial timber sale, in this case the biomass was generated from the entire tree, not merely the tops.

In 2014 poor markets resulted in 789 cords to be chipped. This was counted as round wood and deducted from the allowable harvest, rather than biomass byproduct.

In 2015, hemlock markets were poor, and as a result 972 cords of whole tree hemlock were chipped for biomass.

In 2016, hemlock markets were poor, and as a result 924 cords of whole tree hemlock were chipped for biomass.

³ Biomass in 2017 was also generated from a wildlife habitat project that had very little commercial value. In this case the biomass was generated from the entire tree, not merely the tops.

⁴ Softwood Pulp was composed of a mixture of White Pine and Hemlock pulpwood at the end of deliveries in 2019. For a few weeks, that was the only way to sell Hemlock pulpwood.

Other Forest Products

DLLT, as part of its community forest management, routinely issues permits to local users of forest products, subject to policies and procedures approved by the DLLT Board of Directors. A voluntary survey is mailed to each permit holder the following year to keep track of products harvested.

In 2020, DLLT issued permits for gravel, firewood, “tips” for wreaths, trees and craft wood.

Gravel: 11 permits issued, (11 reported) 410 cubic yards sold

Firewood: 58 permits issued (29 reported), 33 cords reported

Tipping: 16 permits issued (4 reported), 1,325 lbs reported

Trees: 1 permits issued (0 reported) 10 trees used

Craft wood: 3 permits issued, (3 reported) 7 trees used

Unanticipated Removal or Loss

DLLT staff and forestry contractors monitor the forest for unanticipated loss due to insects, disease, wind, fire, excessive browsing by animals, and timber theft during routine management operations. DLLT also uses reports from members and others who use the forest to keep informed of changes in the forest.

No unanticipated losses occurred in 2020.

Regeneration

DLLT staff, board members, and forestry contractors monitor forest harvest areas to determine if regeneration is occurring as anticipated and intended in forest harvest plans. Qualitative or quantitative inspections generally will occur within three years of harvests intended to encourage regeneration.

2015 Regeneration and Monitoring Statement

Background: As specified in the management plan, 2015 harvest regeneration monitoring was conducted after 5 growing seasons, in February 2021. This gives forest managers an opportunity to see any other progress in the development of post harvest stands. If undesirable conditions are observed, they were noted to avoid those conditions in the future.

Results:

Area 1 South of 4th Lake Road near Mile 10 (Areas 1-5 in T5ND)

- **Light to moderate thinning:** This mixedwood area has residual basal areas ranging from 40 – 60 sq.ft. per acre. There is a good stocking of advance regeneration which pre-existed the harvest, with raspberry dominating the skid trails. Forest health is good.

Monitoring Summary Report

Area 2 West of road to Buck Knoll

- **Liberation Patches:** This softwood area has scattered mature trees and much advanced softwood regeneration often overstocked. There is evidence of moose browsing of hardwood and raspberries are plentiful in some skid trails. Merchantable residual basal area ranges from 20 to 60 sq.ft.per acre.

Area 3 North of 4th Lake Road – near mile 10.5

- **Light and Moderate Thinning:** This softwood area has residual basal areas ranging from 40 – 70 sq.ft. per acre including some tolerant hardwood. There is a good to overstocked advanced mixed regeneration which pre-existed the harvest, with raspberry dominating the skid trails. Forest health is good. Found a Hemlock bumper tree that died since harvest. Check other harvests for similar instances.

Area 4 Just NW of Area 3.

- **Liberation Patches:** Thick advanced mixed species regeneration. Only 10 – 20 sq.ft. of residual basal area which conforms to the prescription. There is plenty of evidence of moose browsing of hardwood and softwood species. Precommercial thinning is a possibility, although many of the softwood are under 5 feet tall.

Area 5 Between 4th Lake Road and North boundary near mile 11

- **Light and Moderate Thinnings:** Mixedwood area with scattered Sugar Maple. Basal areas from 40-80 sq.ft. Advance regeneration is from good to overstocked. Moose are browsing most species. Forest health is good.

Note: Other 2015 operating areas described in T6ND were either not operated or checked before.

Focus Species Habitat Management Activities

Management for specific “focus species” is used to benefit species of interest to the local community and to provide habitat for the full range of wildlife species found on the forest. The management plan sets out specific management activities for these species.

DLLT Self-Designated Deer Management Areas

DLLT has a significant, long-term goal of restoring and expanding deer wintering areas, the availability of which is a limiting factor for local deer survival. Management activities include both building the area of mature forest softwood cover through partial harvesting in historic deer wintering areas (primary and secondary cover), harvesting to create openings that will produce browse and regenerate the forest to ensure a steady supply of future winter cover, and seeding landings to create summer food for deer and other species.

Deer Wintering Area Management Activities							
Habitat and Activity	2008- 2014 (ac)	2015 (ac)	2016 (ac)	2017 (ac)	2018 (ac)	2019 (ac)	2020 (ac)
Partial harvests (selection, initial	147	0	0	30	21	30	210

Monitoring Summary Report

shelterwood, and intermediate harvests) ²							
Regeneration harvest openings (patch-cut, overstory removal, and clearcut ²	49	0	0	0	0	110	0
Herbaceous seeding ¹	0	0	0	0	0	1	0
Management consistent with DWA 5-year operations plan	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Other Monitoring:							

During the Summer of 2020, timber harvesting was conducted near Spring Cove, part of our Deer Management Areas. The harvest in Primary and Secondary cover was a light thinning to preserve cover.

Snowshoe Hare

The best snowshoe hare habitat is created by even-aged regeneration harvests in softwood-cover. The “regeneration harvest openings” for deer wintering area management are also used to monitor the amount of snowshoe hare habitat created. Snowshoe hare habitat is consistent with nearly all DLLT’s shade tolerant conifer management as well.

Grouse and Woodcock

Grouse and woodcock management is based on creating a number of patches of different age classes in aspen and birch stands. The following monitoring elements have been included to track progress toward objectives outlined in the management plan.

Grouse and Woodcock Management

¹ DLLT also keeps track of species and location of species used in herbaceous seeding.
² Harvest near Spring Cove in T5 and T6ND – partial cuts in DMAs in 2020

Monitoring Summary Report

Annual Monitoring Element	2008-2014	2015	2016	2017	2018	2019	2020
Number of grouse/woodcock unit plans developed	Seven	None	None	None	One	None	None
Cumulative number of units designated for active management	Eight	Eight	Eight	Nine	Nine	Nine	Nine
Total number of acres harvested (clearcut or overstory removal) across all management unit blocks	97.9	42	0	80	0	48	0
Number of acres of herbaceous seeding	Landings Seeded Summer 2013	Landings Seeded Summer 2014	Landings Seeded Summer 2015	Landings Seeded Spring 2016	None	Landings seeded Summer 2019	Landings seeded Fall 2020

* In the 2008 summer harvest, a set of seven patch cut harvest blocks in a poplar-birch fire origin stand on the south side of Burroughs Brook on the Farm Cove peninsula were created to provide early-successional habitat, including habitat for Grouse and Woodcock and browse for deer and moose. Average block size was 0.56 acres, with just under 4 acres harvested in total. A complete plan for this grouse/woodcock unit has not yet been developed, but the harvest plan calls for a 10-year re-entry to harvest new ½ acre patches adjacent to the patches harvested in 2008.

* In the 2009 summer harvest, a set of 3 patch cut harvest blocks in a 28 acre poplar-birch fire origin stand east of Burroughs Brook on the Farm Cove Mountain Road were created to provide early successional habitat, including habitat for Grouse and Woodcock and browse for deer and moose. The average patch size was 0.3 acres, with under 1.25 acres harvested in total.

*In the 2010 winter harvest, 130 acres of designated Grouse/Woodcock habitat were included on the Farm Cove Peninsula. In the 2010 summer harvest, there were 22 acres of designated Grouse/Woodcock habitat included. Upon closer inspection of designated grouse areas, it was determined areas were either un-merchantable, or were currently unsuitable for grouse and woodcock management.

*In the 2011 winter harvest, 9 acres were harvested in 8 patches of between 2 and 0.5 acres each. These areas were dominated by mixed intolerant hardwoods, with some scattered hemlock and pine. This occurred on the Daugherty ridge road. In the summer of 2011, 4 clearcuts were created ranging between 3 and 6 acres in size. These patches were irregular and occurred mostly in intolerant hardwoods, with some mixed spruce and hemlock. This occurred on the Dobsis dam road.

* In the 2012 summer harvest, a plan encompassing roughly 100 acres was developed to provide early successional habitat on a rotational basis. In 2012 approximately 16 acres were clearcut in 2.5 and 4.5 acre circles along the 4th lake road between .5 and 1 mile markers.

* In the 2013 the harvest unit established in 2012 was completed, totaling 37 acres in the entire harvest unit. In addition, a new harvest unit was established on Norway point, east of farm cove. This consisted of a 40 acre stand where approximately half the area was clearcut in the form of 2 acre rectangles, in a checkerboard fashion, totaling approximately 22 acres.

Monitoring Summary Report

* In 2015 the Belden Brook grouse and woodcock management plan was continued for its second entry. A total of 42 acres was created. Five 3 acre clearcuts were included in this project for the NRCS, qualifying us to continue with the CSP program. Other patches were generally larger, but within Cat. I status.

*In 2017, the first entry of patch-cuts was implemented on the Coca Cola Highway. A total of eight patches totaling 35 acres was cut. 11 patches were cut on Sonny's Lane, totaling approx. 45 acres.

*In 2019, the second entry of the first phase patch-cuts was completed on the Coca Cola Highway. 11 patches totaled approximately 48 acres.

The balance of aspen-birch age classes on the entire forest is also monitored periodically as cover type maps are updated (see Section III).

Black Bear

Black bear habitat management is accomplished through our creation and maintenance of young-forest openings by implementing the grouse/woodcock and deer wintering area management plans and implementation of the hard mast guidelines during harvest operations. These activities are monitored, and we conduct no separate monitoring of black bear habitat conditions or management.

Riparian Zone Management

Harvest and other operations monitoring forms are used to gather information on harvest activities within riparian management areas. A summary of problems identified (e.g., unsatisfactory performance relative to management plan guidelines or site-specific plans) and steps taken to correct problems described below.

Year	Unsatisfactory Implementation of RMZ Guidelines and Action Taken				
	Trout/Salmon	Beaver	Lake	Other Stream	Vernal Pool
2008-2015	No problems observed	No problems observed	No problems observed	No problems observed	No problems observed
	Action:				
2016	No problems observed	No problems observed	No problems observed	No problems observed	No problems observed
	Action:				
2017	No problems observed	No problems observed	No problems observed	No problems observed	No problems observed
	Action:				
2018	No problems observed	No problems observed	No problems observed	No problems observed	No problems observed
	Action:				
2019	No problems observed	No problems observed	No problems observed	No problems observed	No problems observed
	Action:				
2020	No problems observed	No problems observed	No problems observed	No problems observed	No problems observed
	Action:				

Beaver

Habitats modified by beaver activity have been shown to be beneficial to a wide range of wildlife, including waterfowl, wading birds, migratory songbirds, and moose. Other mammals are such as deer and bear are attracted to the early flush of nutritious vegetation in spring. Recent studies from the Moosehead lake region have found that rusty blackbirds (a declining species listed as Special Concern in Maine) were strongly associated with beaver-impounded wetlands, and olive-sided flycatcher (also Special Concern) was also found in these areas (Pelletier and Arsenault 2007). Maine has a long history of habitat management guidance that recognizes the benefits of maintaining beaver activity in the landscape, including Deifenbach et. al 2008, Foss 1999, and Bryan 2007. Only one stream in the DLCF, Burroughs Brook, has been designated as a priority beaver habitat in the Focus Species Addendum. Burroughs Brook is a slow moving stream with historic beaver use and forest cover that is less dense than that on streams with priority for brook trout and Atlantic salmon.

Brook Trout / Atlantic Salmon

No fish passage improvements were installed in 2020.

Exotic and Invasive Plants

DLLT monitors the use of exotic (non-native) species to ensure that they do not become invasive. Currently DLLT's use of exotic species is limited to planting non-invasive grasses and legumes for wildlife habitat improvement. In addition, DLLT checks for the presence of known invasive plants that may be present in the area. A new tool called the Maine Invasive Plants Field Guide is now in use by DLLT staff.

Monitoring Summary Report

Exotic and Invasive Plants						
	2008- 2015	2016	2017	2018	2019	2020
Wildlife Plantings						
Number of sites planted	harvest landings from 22 seasons and excavated ditches,	Summer 2015 and winter 2016 harvest landings and excavated ditches	Summer 2016 and winter 2017 harvest landings and excavated ditches		Summer 2019 harvest landings and excavated ditches	Summer 2020 harvest landings and excavated ditches
Species	Conserv. Mix (Contains non-native grasses & legumes)	Conserv. Mix (Contains non-native grasses & legumes)	Conserv. Mix (Contains non-native grasses & legumes)	Conserv. Mix (Contains non-native grasses & legumes)	Conserv. Mix (Contains non-native grasses & legumes)	Conserv. Mix (Contains non-native grasses & legumes) ³
Estimated total area planted	3 acres	?	2 acres		2 acres	2.1 acres ⁽³⁾
Seed mix does not contain species on Maine's list of invasive plants (Y/N)	Y	Y	Y	Y	Y	Y
Location identified in GIS (Y/N)	Y	Y (general location of harvest areas and roads)	Y (general location of harvest areas and roads)	Y (general location of harvest areas and roads)	Y (general location of harvest areas and roads)	Y (general location of harvest areas and roads)
Number of sample sites checked for undesirable spread	Twenty seven (Earlier Plantings)	Three (earlier plantings)	Three (earlier plantings)	None	None	None
Undesirable spread noted?	No	No	No	No	No	No
Invasive Plants						
All harvest sites checked?	Yes, During routine operations and tour	Yes, During routine operations and tour	Yes, During routine operations and tour	Yes, During routine operations and tour	Yes, During routine operations and tour	Yes, During routine operations and tour
Species found? ^{1,2}	No	No	No	No	No	No

Monitoring Summary Report

¹ Describe severity or impacts of any invasive species or exotic species and develop an action plan if management is feasible and warranted.

² Staff received training and procured the new "Maine Invasive Plants Field Guide".

³ Staff planted pollinator seed mix, 0.1 acre at Billy Brown Field.

Harvest Impacts

Harvest sites and road improvement projects are monitored by DLLT’s Forest Manager, Executive Director, and Board of Directors to ensure compliance with applicable laws and Best Management Practices designed to protect soil and water quality. Harvest operations are also monitored to ensure that operations comply with silvicultural prescriptions, damage to standing timber and regeneration is minimized, sensitive sites are protected, and site-specific wildlife practices and objectives are being met.

	Compliance with Harvest Guidelines					
Monitoring element/guideline	2008-2015	2016	2017	2018	2019	2020
Hard Mast referenced in harvest plans for applicable stands	S	S	NA	S	S	S
Satisfactory execution of hard mast guidelines during harvest	S	S	S	S	S	S
Wildlife trees and downed logs	S	S	S	S	S	S
Retention patches	S	S	NA	NA	NA	NA
Wildlife Trees retention Patch: quantitative sample of selected harvest blocks (number of blocks, performance)	S	NA	S	NA	NA	NA
Riparian and Lakeshores: applicable guidelines referenced in management plans	S	S	NA	S	S	S
Riparian and Lakeshores: applicable guidelines and BMPS implemented	S	S	S	NA	S	S
BMPs beyond riparian and lakeshore zones	S	S	S	S	S	S
# vernal pools known prior to harvest plan:	0	0	0	1	0	0
# new vernal pools identified (1	0	0	0	0	2	0
Vernal pools identified in harvest plans and guidelines implemented during harvest	NA	NA	NA	S	S	NA

S – Satisfactory

U - Unsatisfactory, problem ongoing (describe below)

U/S – Unsatisfactory, problem corrected (describe below)

Except where noted above, all harvests are monitored for all element
(1 pending verification by biologist

Monitoring Summary Report

Unsatisfactory Harvest Conditions: Identification and Resolution

During 2020, harvest conditions were generally satisfactory in terms of both silvicultural and ecological objectives. The harvest contractor has continued to demonstrate a strong understanding of DLLT goals and objectives.

Road Monitoring

Road Monitoring Summary	
YEAR	Roads Inspected, Problems Identified and Corrected
2008	4 th Lake Rd: Entire road monitored; previously approved maintenance project completed, including installation of 10 culverts, and ditching and re-shaping on portions of 7 miles of the road, and routine grading occurred. Brushing of the road way was completed in 2007. Installation of a new bottomless arch culvert at Rolfe Brook to improve aquatic habitat and fish passage completed. Additional ditching, culvert, and graveling work is planned for 2009 and beyond.
	Farm Cove Dam Rd.: entire road monitored, brushing completed; installation of new bottomless arch culvert at Scott Brook to improve aquatic habitat and fish passage and two new nearby cross-drain culverts completed
	Farm Cove Mountain Rd: road north to Burroughs Brook monitored; ditching and surface maintenance completed as needed to support harvest activities.
	Dobsis Dam Rd: entire road monitored; surface condition poor and limited surface erosion occurring; no substantial watershed impacts but maintenance improvements recommended for recreational use as funds available. Roadside brushing was completed in 2007.
2009	4 th Lake Rd: Entire road monitored; routine seasonal gradings conducted, culvert replaced at mile 8.6.
	Farm Cove Dam Rd.: Entire road monitored; culvert replaced at mile 2.8
	Farm Cove Mountain Rd: road north to Burroughs Brook monitored; ditching and surface maintenance completed as needed to support harvest activities.
	Dobsis Dam Rd: entire road monitored; surface condition poor and limited surface erosion occurring; no substantial watershed impacts but maintenance improvements recommended for recreational use as funds available.
	Third Lake Rd: on Wabassus Tract acquired 12/08: entire road monitored; surface condition poor and limited surface erosion occurring; no substantial immediate watershed impacts but maintenance improvements recommended. Arch culvert installed at Wabassus tributary stream (see "Brook Trout" above). Major erosion risk exists on section of road below Wabassus Mt with highly eroded ditch that lacks functional cross drains; drainage restoration project planned for 2010.
	43-00-0 / Little River Rd: on Wabassus Tract acquired 12/08: entire road monitored; surface condition poor and limited surface erosion occurring; no substantial immediate watershed impacts but maintenance improvements recommended. Major portions lack adequate drainage ditches or cross drains and are at risk for erosion, surface extremely rough. Restoration project planned for 2010
	42-00-0 / Little River Rd: on Wabassus Tract acquired 12/08: entire road monitored; surface condition poor and limited surface erosion occurring; no substantial immediate watershed impacts but maintenance improvements recommended. Lacks adequate drainage ditches or cross drains and is at risk for erosion, surface extremely rough. Restoration project planned for 2010
2010	Wabassus Mt Rd: on Wabassus Tract acquired 12/08: entire road monitored; surface condition poor and limited surface erosion occurring; no substantial immediate watershed impacts. Lower priority for restoration.
	4 th Lake Rd: Entire road monitored; routine seasonal gradings conducted
	42-00-0 and 43-00-0 Rds: Drainage and surface restoration projects completed, including culvert installations
	88 Rd: arch culvert installed at North Br; see "brook trout" above.
2011	Wabassus Mt Rd: arch culvert installed at North Br; see "brook trout" above.
	4 th Lake Rd & Little River Rd: Entire roads monitored; routine seasonal gradings conducted

Monitoring Summary Report

	3rd Lake Rd (60-00-0): Drainage and surface restoration project completed on northern portion of road, including culvert installations
2012	4 th Lake Rd & Little River Rd: Entire roads monitored; routine seasonal gradings conducted Wabassus Mt. Rd, portion of 3 rd Lake Rd, portion of Dobsis Dam Rd, drainage and surface restoration project completed, including culvert installations; small culvert installation on ATV trail near Billy Brown field
2013	4 th Lake Rd & Little River Rd: Entire roads monitored; routine seasonal gradings conducted Planned gradings completed on 3 rd Lake Ridge Rd, Dobsis Dam Rd, Farm Cove Dam Rd, Wabassus Mt Rd. Drainage and surface restoration projects completed on portions of 4 th Lake Rd, 4 th Lake Landing Rd, Belden Brook Rd, 3 rd Lake Ridge Rd, and Farm Cove Dam Rd, including culvert installations; approx. 10.5 miles of roadside brushing completed
2014	4 th Lake Rd & Little River Rd: Entire roads monitored; routine seasonal gradings conducted Planned gradings completed on 3 rd Lake Ridge Rd, Dobsis Dam Rd, Farm Cove Dam Rd, Wabassus Mt Rd Drainage and surface restoration projects completed on portions of 4 th Lake Rd, including culvert installations.
2015	4 th Lake Rd & Little River Rd: Entire roads monitored; routine seasonal gradings conducted. 4 culverts were also installed Planned gradings completed on farm cove dam rd, 3 rd lake Ridge road, 3 rd lake outlet road, and a portion of the Beldon brook rd. Gravel was screened for the first time in the Elsemore pit.
2016	4 th Lake Rd & Little River Rd: Entire roads monitored; routine seasonal gradings conducted 4 arch culvert installations, 1 decommissioning 6 drainage culvert installations and 9 miles brushing on Amazon Road
2017	4 th Lake Rd, Little River Rd, Amazon Rd: Entire roads monitored; routine seasonal gradings conducted 1 bridge installation, 3 decommissionings ¼ mile ditching 1 gate installation
2018	4 th Lake Rd., Little River Road, Amazon Rd.: Entire roads monitored; routine seasonal gradings conducted No bridge installations
2019	4 th Lake Road, Little River Road, Amazon Road: Entire roads monitored; Summer grading conducted. 1 arch installation, 40' portable bridge installed twice for harvesting. Gravel screened and stockpiled at Elsemore pit. Approx. 4 miles of brushing done on Wabassus and 3 rd Lake Ridge Road. 6 culverts replaced.
2020	Seven gates were installed to protect main roads from spring damage. 4 th Lake Road, Little River Road, Amazon Road: Entire roads monitored; Summer grading conducted. Gravel screened from the Chet's Camp Road pit and used on 3.7 miles of resurfacing of the Amazon Road. 3.9 miles of ditching on the Amazon Road. Reshaped 4 hills on the Little River Road. Fall grading of 7 miles of the 4 th Lake Road. 6.6 miles of roadside brushing.

Pesticides and Biological Control Agents

DLLT does not currently use pesticides or biological control agents. If in the future a need to use the agents arises, DLLT will prepare evaluate the risks, prepare appropriate application plans, and monitor use in accordance with the Downeast Lakes Community Forest Management Plan, the conservation easement, and Maine law and Forest Stewardship Council certification standards.

Social and Economic Monitoring

Social and Economic Monitoring						
Element	2008-2015 (Avg)	2016	2017	2018	2019	2020
Total volume of wood harvested (cords)	6236	6032	6696	11157	12462	9605
Number of permanent DLLT employees	3	4	4	4	5	5
Number of temporary DLLT employees	0	2	2	4	3	3
Number of contractor and subcontractor employees	Appr. 13*	Appr. 13*	Appr. 10*	Appr*. 10	Appr*. 10	Appr*.10

* contractors and subcontractor employees include foreman, operators, truckers, and other employees of harvest contractor Davis Forestry Products; only harvest-related contractor employees are included here.

DLLT’s board members and staff, and public meetings attended or hosted by DLLT are the major means by which DLLT monitors the public reaction to management.

2020- In the past year, DLLT has heard many positive comments regarding the Trust’s road management, conservation of lands, and efforts to preserve the lakeshores. Specific comments included positive reactions about the resurfacing of the Amazon Road. We also received positive comments regarding the expansion of the hiking trail system.

Common Loon Monitoring

In addition to monitoring activities directly related to management of the Downeast Lakes Community Forest (formerly the Farm Cove Community Forest) described in this report, DLLT has had a program of monitoring common loon productivity on lakes throughout the region, a project to establish a baseline set of data on loon reproduction that began in 2001 in cooperation with the U.S. Fish and Wildlife Service and Biodiversity Research Institute, and also involved Maine Audubon in 2011. This monitoring effort concluded in 2011. An executive summary of

Monitoring Summary Report

the loon monitoring report is available upon request.

III. Periodic Forest Monitoring Data

Because the following data are gathered periodically (for example, every 5-10 years), this section of the report will be only updated as new data become available.

Forest Inventory

Standing Timber

Forest inventory is the basis of good forest management. The following is a summary of the data that have been collected on the forest. A forest-wide inventory took place in 2000 on the original Farm Cove Property. An inventory of the Wabassus Lake Tract was completed in 2008. In the fall of 2014, aerial photos and stand type maps were developed.

New inventory points were completed in 2010 on the original Farm Cove Community Forest, exclusive of the Wabassus Lake Tract. After processing the cruise data collected by Fountains Forestry, the following results were compiled using the 7 Islands inventory program, MBG tools. The following table contains current cruise information (2010) and past cruise data and estimates.

Summary	Entire Ownership (excl Wabassus)	Without LSMA or Eco Reserve	Without Eco Reserve	LSMA Only	Ecological Reserve Only
Cords per Acre 2003	17	?	?	?	?
Estimated Cords per Acre 2008	18.1	17.6	17.5	17	21
Cords per Acre 2010	18.4	17.8	18.1	19.3	19.7

In 2015, this inventory information was updated in 2014 with re-measured post-harvest plots, and “grown” plot information received from Finite Carbon as part of our Carbon monitoring program. This data was processed using LMS (Landscape Management Systems) and formulas developed by the Forest Service, to update the inventory once again. The results were not surprising, there is a clear annual net gain. Harvest rates are approximately half of growth. More inventory information can be delivered upon request.

Based on the most recent ***round wood*** cruise data, the following has occurred:

- Farm Cove Tract, 2005-2015
 - The average net (growth-harvest) increase in stocking from 2005 to 2015 was 0.35 cords per acre per year.
 - The Average Gross Growth (excluding harvest) was 0.63 cords per acre per year.
- Wabassus Tract, 2007-2014

Monitoring Summary Report

- The average net (growth-harvest) increase in stocking from 2007-2015 was 0.31 cords per acre per year.
- The Average Gross Growth (excluding harvest) was 0.47 cords per acre per year.
- West Grand Lake Tract
 - The average net (growth-harvest) increase in stocking from 2009-2014 is unknown. The property was not in the possession of DLLT until July of 2016.
 - The Average Gross Growth (excluding harvest) is unknown. The property was not in the possession of DLLT until July of 2016.

*Numbers are based on roundwood only.

In 2015 most of the Farm Cove Community Forest was re-inventoried with fixed 1/15 acre plots as part of our FC Carbon project. This new inventory, although conducted on the same plot location, came out with an additional 76,312 cords, averaging out to an additional 3.6 cords per acre (25.6 cords/acre). With the addition of the West Grand Lake Tract, our Annual Allowable Cut (AAC) has increased to 12,000 cords per acre per year.


Changes in Habitat Conditions

The Downeast Lakes Community Forest is managed for a range of forest types and ages to provide diverse and abundant habitat for wildlife species of interest to the local community. Aerial photographs and cover type maps are used to assess forest habitat conditions for most species. To help manage the forest, the management plan has divided the forest into the following management units.

LSMA	North of the 4 th Lake Ecological Reserve including Beldon Brook Area and southwest of Third Lake Ridge Road
Dark Cove:	South of Dobsis Dam to Fourth Lake Road and West to Buck Knoll
McLellan Cove:	North of West Grand Lake
Farm Cove:	South of West Grand Lake and east of the Wabassus-Pocumcus thoroughfare. Includes 30 acres on Kitchen Cove Point
Wabassus Lake:	T43, area south of Fourth Lake Road
Bonney Brook:	Areas below the Milford Road, and along the Bonney brook Rd
Amazon:	Area to the North and South of the West Grand Lake Ecological reserve
Amazon-Musquash Ecological Reserve:	East side of Amazon Road and south of Otter Brook Road. Also West side of Amazon Road north of Otter Brook Road

Monitoring Summary Report

The graph below represents forest habitat conditions as of 2015 summarized from the cover type data. The next update of the cover type maps and data is expected in or before 2025.



Deer Management Areas

Long-term monitoring of deer wintering areas is based on the percent of mapped primary and secondary cover in mapped DMAs. Based on aerial imagery from 2014, DLLT has identified eight management areas totaling 19,836 acres. The objective is to have at least 25% of each DMA in primary cover and at least 50% in primary and secondary cover combined. Due to heavy harvesting under previous ownership, none of the areas meet the state DWA cover criteria.

Deer Wintering Areas														
DMA	Primary		Secondary		Primary + Secondary		Future Secondary		Never Cover		NA		Total	Goals Met?
	%	ac	%	ac	%	ac	%	ac	%	ac	%	ac	ac	Yes/No
Amazon	8%	544	23%	1649	31%	2193	48%	3396	3%	212	18%	1255	7055	No
Belden Bk	43%	696	20%	335	63%	1031	23%	373	11%	177	3%	56	1637	Yes
Burroughs Bk	36%	323	33%	292	69%	615	21%	187	8%	73	2%	20	895	Yes
Bonney Bk	11%	202	4%	74	16%	276	43%	770	12%	215	13%	238	1775	No
Grand Lake Bk	22%	664	13%	398	36%	1062	58%	1727	0%	0	7%	201	2990	No
Hayes Bk	30%	639	24%	515	53%	1154	41%	887	3%	54	3%	63	2159	Yes
Wabassus	18%	424	13%	299	31%	723	53%	1232	12%	276	5%	106	2338	No
Whitney Cove	34%	339	28%	281	63%	620	35%	343	2%	18	1%	5	987	Yes
Grand Total	20%	3831	19%	3843	39%	7674	45%	8915	5%	1025	10%	1944	19836	No

Monitoring Summary Report

*Total cover includes all primary, secondary, and non-cover areas.

** GL Brook was increased to match acreage of marten management area

Section II includes a summary of annual management activities in DMAs.

American Marten

The management plan for American marten is based on maintaining large patches (over 1,200 acres) of mature forest. Monitoring is based on periodic inventories and cover type maps (i.e., every 10 years) to quantify habitat conditions.

Marten Habitat by Management Unit

Unit	Total Area	Potential Habitat	Primary Habitat		Secondary Habitat		Current Habitat	
	ac	ac	ac	%	ac	%	ac	%
Amazon	7055	5818	4115	70%	696	12%	4810	82%
Belden Bk	1637	1581	1074	68%	466	29%	1540	97%
Bonney BK	1733	1496	945	63%	476	32%	1422	95%
Burroughs Bk	895	837	594	71%	240	29%	834	100%
Grand Lake Bk	2990	2769	1884	68%	736	27%	2620	95%
Hayes Bk	2159	2024	1542	76%	313	15%	1855	92%
Wabassus	2338	2195	1435	65%	259	12%	1693	77%
Whitney Cove	987	967	515	53%	414	43%	929	96%
Grand Total	19794	17687	12104	68%	3600	20%	15703	89%

Grouse and Woodcock

Long term potential high-value grouse and woodcock habitat is indicated by the total area and balance of development stages in the aspen-birch forest type. This is only a portion of the total area of habitat, because grouse will also be found in young and intermediate-aged northern hardwood and hardwood-dominated mixed forests.

2005 Grouse and Woodcock Habitat Conditions						
Management Unit	Focus Species Development Stage (ac)				Aspen-Birch Total	All Types Total
	Early Successional R/S	Early Successional S/SP	Intermediate	Mature		
Dark Cove (exclusive of the Ecological Reserve and LSMA)	28	444	76	14	562	4,546
Whitney Cove		46	187	60	293	2,703
Farm Cove		206	422		629	11,992
Aspen-Birch Total	28	697	685	74	1,484	19,240
Total Forest Acres						25,369

2015 Grouse and Woodcock Habitat Conditions						
Management Unit	Focus Species Development Stage (ac)				Aspen-Birch Total	All Types Total
	Early Successional R/S	Early Successional S/SP	Intermediate	Mature		
Amazon	8	119	157	6	290	10,212
Bonney Brook		64	505	270	839	4,549
Dark Cove		121	57		178	4,746
Farm Cove	12	76	510		598	12,911
4 th Lake Ecological Reserve		62	372		434	3,696
LSMA	61	59	438		558	3,568
McLellan Cove			148		148	2,408
Wabassus		15	209	7	232	6,628
A-M Ecological Reserve	7	122	242		371	7,092
Aspen-Birch Total	88	638	2638	283	3648	
Total Forest Acres						55,930

When examining the updated data, it's clear that the acres of aspen birch forest type is decreasing over time. The baseline (2005) data for Dark Cove, McLellan Cove, and Farm Cove had a total of 1,484 acres of Aspen-Birch forest type. That number has dropped to 924. This is due to forest succession and harvesting practices. Many aspen birch stands have developed significant shade tolerant conifer cohorts, making it difficult to justify retaining cover type.

DLT is considering several options in preserving the presence of the forest type.

Black-throated Blue Warbler/Mature Hardwood Forest

Black-throated blue warbler is the focal species for older intermediate and mature northern hardwood forest. In 2005 about 60% of this forest type in the DLLT FCCF as a whole was in the early successional stage and 6% was in the mature stage. The objective was to increase mature northern hardwoods to 15% of the total northern hardwood area by 2015 and 30% by 2025.

With the current updated cover type maps, changes were observed. Currently, 36% of northern hardwood forest is in the early successional stage. This shows an increase in stand development. We also increased TOTAL northern hardwood acreage on the DLCF. With the new total acreage, we have actually seen a slight decrease in mature hardwood acreage, in percent cover. If the original acreage is used to calculate current mature cover, than we have actually increased to 7% mature cover.

It is believed that the increase in total acres in northern hardwood cover is due to forest succession, or possibly some minor mis-typing in forest types in 2005. It is also believed that the growth rates required to achieve mature northern hardwood status is may have been underestimated.

An 8 acre Crop Tree Release was initiated near GLS in T6ND. More than half of these acres remain to be treated. A few acres in the front are complete and a kiosk explaining the treatment was installed near the main road.

Hard Mast Management

Long-term plans for hard mast include experimental planting of American chestnut and red oak by 2012. These plots and chestnuts planted prior to 2007 will be monitored.

Year	Number of plots	Year Planted	Type of Planting	Year Monitored	Results
Planted Prior to 2008	8	2006	Seedlings	2008	Qualitative inspection only; surviving seedlings appear healthy and have received only moderate browse pressure; survival appears better away from raspberry vines in old wood yards
2008					
2009					
2010					
2011					
2012					
2013					
2014					The planting of Chestnut is being reconsidered, but in a drastically different manor than before. Any future plantings will likely focus on Red oak.

Monitoring Summary Report

2015					
2016					
2017					
2018					
2019					
2020					

Rare Species, Natural Communities, and other Special Habitats

During 2002-2003 DLLT contracted with Dr. Norm Famous and Janet McMahon to inventory the anticipated DLLT acquisition lands for the presence of rare, threatened, or endangered wildlife and plant species. The final report and recommendations were completed in August of 2007. Additional information, including a list of rare species that could potentially be observed on the Community Forest, was requested and received from the Maine Natural Areas Program, and is summarized in the Farm Cove Community Forest Management Plan (2008). The Wabassus tract was acquired in 2009, and a focus species plan was developed in 2010. With the acquisition of the West Grand Lake Tract in 2016, DLLT decided to combine all management into a single document, Downeast Lakes Land Trust Forest Management Plan (2015). DLLT's approach is to protect species by protecting their habitat, including areas designated as special management areas, late-successional forest, and ecological reserve. Monitoring for general conditions or unintended adverse impacts occurs primarily during forest harvest operations planning and implementation when harvests occur near or in special management areas.

Literature Cited

Bryan, R.R. 2007. Focus Species Forestry: A Guide to Integrating Timber and Biodiversity Management in Maine. Maine Audubon, Falmouth.

Deifenback, D.R, S.J. Lovett, and R.B. Owen, Jr. Beaver and wetlands. 1988. P. 38-41 in Elliott, C.A., ed., A Forester's guide to Managing Wildlife Habits in Maine. University of Maine Cooperative Extension Service and Maine Chapter of the Wildlife Society.

Foss, C.R. 1999. Beaver influenced wetlands. P. 61-64 in Elliot, C.A., ed. Biodiversity in the Forests of Maine: Guidelines for Land Management. UMCE Bulletin 7147. University of Maine Coop Extension, Orono.

Pelletier, S.P. and M.P. Arsenault. 2007. Pre-filed direct testimony on Behalf of Plum creek Timber Company in support of the Moosehead Region Concept Plan.

ftp://ftp.state.me.us/outgoing/PlumCreek/ReceivedFromPlumCreek/2007-08-31%20PCFilings/08-31-2007%20PrefiledDirectTestimony/Volume04/2007-08-31_PLUMCREEK_WoodlotTestimony.pdf